
Supplementary Material: Efficient Algorithms for Robust One-bit Compressive Sensing

Lun Zhang

ZHANGLJ@LAMDA.NJU.EDU.CN

National Key Laboratory for Novel Software Technology, Nanjing University, Nanjing 210023, China

Jinfeng Yin

JINFENGY@US.IBM.COM

IBM Thomas J. Watson Research Center, Yorktown Heights, NY 10598, USA

Rongjin

RONGJIN@CSE.MSU.EDU

Department of Computer Science and Engineering, Michigan State University, East Lansing, MI 48824, USA

A Proof of Lemma 6

We consider the following general optimization problem

$$\min_{\|\mathbf{x}\|_2 \leq 1} -\mathbf{x}^\top \mathbf{y} + \gamma \|\mathbf{x}\|_1. \quad (15)$$

Before we proceed, we need the following lemma.

Lemma 6 *The solution to the optimization problem*

$$\min_x \frac{1}{2}(x - y)^2 + \gamma|x|$$

is given by

$$P_\gamma(y) = \begin{cases} 0, & \text{if } |y| \leq \gamma; \\ \text{sign}(y)(|y| - \gamma), & \text{otherwise.} \end{cases}$$

where $P_\gamma(\cdot)$ is the soft-thresholding operator defined in (7) (Donoho, 1995).

The proof of Lemma 6 can be found in (Duchi & Singer, 2009). Based on the above lemma, it is easy to verify that

$$\min_x \frac{1}{2}(x - y)^2 + \gamma|x| = \begin{cases} \frac{y^2}{2}, & \text{if } |y| \leq \gamma; \\ \gamma|y| - \frac{\gamma^2}{2}, & \text{otherwise.} \end{cases} \quad (16)$$

First, we consider the case $\|y\|_\infty \leq \gamma$. Then, it is easy to verify that

$$\mathbf{0} \in \underset{\mathbf{x}}{\text{argmin}} -\mathbf{x}^\top \mathbf{y} + \gamma \|\mathbf{x}\|_1.$$

Since $\|\mathbf{0}\|_2 \leq 1$, $\mathbf{0}$ is also an optimal solution to (15).

Next, we consider the case $\|y\|_\infty > \gamma$. Following the standard analysis of convex optimization (Boyd & Vandenberghe, 2004), the Lagrange dual

function $g(\mu)$ of (15) is given by

$$\begin{aligned} g(\mu) &= \min_{\mathbf{x}} -\mathbf{x}^\top \mathbf{y} + \gamma \|\mathbf{x}\|_1 + \mu(\|\mathbf{x}\|_2^2 - 1) \\ &= \min_{\mathbf{x}} 2\mu \left(\frac{1}{2} \left\| \mathbf{x} - \frac{\mathbf{y}}{2\mu} \right\|_2^2 + \frac{\gamma}{2\mu} \|\mathbf{x}\|_1 \right) - \frac{\|\mathbf{y}\|_2^2}{4\mu} - \mu \\ &= 2\mu \left(\sum_i \min_{x_i} \frac{1}{2} \left(x_i - \frac{y_i}{2\mu} \right)^2 + \frac{\gamma}{2\mu} |x_i| \right) - \frac{\|\mathbf{y}\|_2^2}{4\mu} - \mu \\ &\stackrel{(16)}{=} 2\mu \left(\sum_{i:|y_i| \leq \gamma} \frac{y_i^2}{8\mu^2} + \sum_{i:|y_i| > \gamma} \left(\frac{\gamma|y_i|}{4\mu^2} - \frac{\gamma^2}{8\mu^2} \right) \right) \\ &\quad - \frac{\|\mathbf{y}\|_2^2}{4\mu} - \mu \\ &= \sum_{i:|y_i| > \gamma} \left(\frac{\gamma|y_i|}{2\mu} - \frac{\gamma^2}{4\mu} - \frac{y_i^2}{4\mu} \right) - \mu \\ &= - \frac{\sum_{i:|y_i| > \gamma} (|y_i| - \gamma)^2}{4\mu} - \mu = - \frac{\|P_\gamma(\mathbf{y})\|_2^2}{4\mu} - \mu. \end{aligned}$$

So, the Lagrange dual problem is

$$\max_{\mu \geq 0} - \frac{\|P_\gamma(\mathbf{y})\|_2^2}{4\mu} - \mu$$

and the optimal dual solution is

$$\mu_* = \frac{\|P_\gamma(\mathbf{y})\|_2}{2}.$$

Following the standard analysis (Boyd & Vandenberghe, 2004, Section 5.5.5), the optimal primal solution is

$$\begin{aligned} \mathbf{x}_* &= \underset{\mathbf{x}}{\text{argmin}} \frac{1}{2} \left\| \mathbf{x} - \frac{\mathbf{y}}{2\mu_*} \right\|_2^2 + \frac{\gamma}{2\mu_*} \|\mathbf{x}\|_1 \\ &\stackrel{\text{Lemma 6}}{=} \frac{1}{\|P_\gamma(\mathbf{y})\|_2} P_\gamma(\mathbf{y}). \end{aligned}$$

B Proof of Lemma 3

We first consider the case $\text{sign}(\mathbf{x}_k^\top \mathbf{u}) = 1$, i.e.,

$$\mathbf{x}_k^\top \frac{\mathbf{u}}{\|\mathbf{u}\|_2} > \delta_k.$$

Then, we have

$$\begin{aligned} \mathbf{x}_*^\top \frac{\mathbf{u}}{\|\mathbf{u}\|_2} &= \mathbf{x}_k^\top \frac{\mathbf{u}}{\|\mathbf{u}\|_2} + (\mathbf{x}_* - \mathbf{x}_k)^\top \frac{\mathbf{u}}{\|\mathbf{u}\|_2} \\ &> \delta_k - \|\mathbf{x}_* - \mathbf{x}_k\|_2 \stackrel{(10)}{\geq} 0. \end{aligned}$$

Thus,

$$\text{sign}(\mathbf{x}_*^\top \mathbf{u}) = \text{sign}\left(\mathbf{x}_*^\top \frac{\mathbf{u}}{\|\mathbf{u}\|_2}\right) = 1 = \text{sign}(\mathbf{x}_k^\top \mathbf{u}).$$

The case that $\text{sign}(\mathbf{x}_k^\top \mathbf{u}_i^k) = -1$ can be proved in a similar way.

C Proof of Lemma 4

First, we have

$$\mathbf{x}_*^\top \mathbb{E}[\mathbf{u}_i y_i] = \mathbb{E}[y_i \mathbf{x}_*^\top \mathbf{u}_i] \stackrel{(4)}{=} \mathbb{E}[\theta(\mathbf{x}_*^\top \mathbf{u}_i) \mathbf{x}_*^\top \mathbf{u}_i] \stackrel{(5)}{=} \lambda,$$

where we use the fact that for a fixed \mathbf{x}_* , $\mathbf{x}_*^\top \mathbf{u}_i$ can be treated as a standard Gaussian random variable.

Consider any vector $\mathbf{x} \perp \mathbf{x}_*$. Since $\mathbf{x}_*^\top \mathbf{u}_i$ and $\mathbf{x}^\top \mathbf{u}_i$ are two independent Gaussian random variable, y_i is independent from $\mathbf{x}^\top \mathbf{u}_i$. Thus, we have

$$\mathbf{x}^\top \mathbb{E}[\mathbf{u}_i y_i] = \mathbb{E}[y_i \mathbf{x}^\top \mathbf{u}_i] = 0.$$

Then, it is easy to prove Lemma 4 by contradiction.

D Proof of Theorem 3

The proof of Theorem 3 is almost identical to that of Theorem 2. The only difference is that in this case, we have

$$\delta_k = \frac{1}{2^{(k-1)/4}},$$

and the total number of calls to the Oracle is upper bounded by

$$\begin{aligned} & m_1 + 2(K-1)t + 2\sqrt{n} \sum_{k=2}^K \delta_k m_k \\ &= m_1 + 2(K-1)t + 2\sqrt{n} m_1 \sum_{k=2}^K 2^{3(k-1)/4} \\ &\leq 2(K-1)t + (3\sqrt{n} 2^{3K/4} + 1)m_1. \end{aligned}$$

E Proof of Corollary 1

We first consider the case that

$$m \leq 2(K-1)t + (5\sqrt{n} 2^{K/2} + 1)m_1,$$

which implies

$$m = O(2^{K/2} \sqrt{n} m_1) = O(2^{K/2} s \sqrt{n} \log n).$$

Thus,

$$\|\mathbf{x}_{K+1} - \hat{\mathbf{x}}\|_2 = \frac{1}{2^{K/2}} = O\left(\frac{s\sqrt{n} \log n}{m}\right).$$

In the case that

$$m \leq m_1 2^K,$$

we have

$$m = O(2^K m_1) = O(2^K s \log n),$$

and thus,

$$\|\mathbf{x}_{K+1} - \hat{\mathbf{x}}\|_2 = \frac{1}{2^{K/2}} = O\left(\sqrt{\frac{s \log n}{m}}\right).$$

F Proof of Corollary 2

The proof is the same as that for Corollary 1.

G Multiplicative Chernoff Bound

Theorem 4. Let X_1, X_2, \dots, X_n be independent binary random variables with $\Pr[X_i = 1] = p_i$. Denote $S = \sum_{i=1}^n X_i$ and $\mu = \mathbb{E}[S] = \sum_{i=1}^n p_i$. We have (Aguilín & Valiant, 1979)

$$\Pr[S \leq (1 - \epsilon)\mu] \leq \exp\left(-\frac{\epsilon^2}{2}\mu\right), \text{ for } 0 < \epsilon < 1,$$

$$\Pr[S \geq (1 + \epsilon)\mu] \leq \exp\left(-\frac{\epsilon^2}{2 + \epsilon}\mu\right), \text{ for } \epsilon > 0.$$

For the second bound, let $t = \frac{\epsilon^2}{2 + \epsilon}\mu$, which implies $\epsilon = \frac{t + \sqrt{t^2 + 8\mu t}}{2\mu}$. Then, with a probability at least e^{-t} , we have

$$S \leq \left(1 + \frac{t + \sqrt{t^2 + 8\mu t}}{2\mu}\right)\mu \leq 2\mu + 2t.$$